



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

therefore

$$Nb = 4271473 \cdot 522,$$

and

$$2Nb = 8542947 \cdot 044.$$

$$\text{From (5) . . . } P \cdot a = \cdot 957 (134920) (t + 5 \cdot 5) + 3 \cdot 56745 (134920) \\ = 129118 \cdot 44 t + 1191511 \cdot 78 ;$$

$$\text{,, (6) . . . } F \cdot c = \cdot 4975 (134920) (t + 5 \cdot 5) - \cdot 4975 (527854 \cdot 4) \\ = 67122 \cdot 7 t + 106567 \cdot 28 ;$$

$$\text{,, (7) . . . } Q \cdot q = 101 \cdot 22798 t^3 + 186 \times 101 \cdot 22798 t^2.$$

Adding and reducing, the equation for *equilibrium* (3) becomes

$$t^3 + 186 t^2 + 1938 \cdot 6 t - 29373 \cdot 24 = 0,$$

whence $t = 8 \cdot 2$ ft. for *equilibrium*.

And the equation for *stability* (4) becomes

$$t^3 + 186 t^2 + 1938 \cdot 6 t - 71570 = 0,$$

whence $t = 14 \cdot 66$ ft. for *stability*.

In the construction of this particular dome, the values obtained above for t will only apply to four points on the walls which carry the dome ; for instead of being built on a circular “ drum,” this drum is made to cover a square chamber by means of “ pendentives,” which are so arranged as to throw the thrust principally on the angles of the building, where the values of a and c will become very great, and it is only at the centre of each of the four sides that the above equations can be employed to compare the thrust of the dome with the strength of the walls. At these four points the walls are about 11 ft. thick, so as to be considerably more than sufficient to produce *equilibrium* ; and the *coefficient of stability* at these points is 1·374 instead of 2. Had the dome therefore been built on a circular drum of 11 feet thickness, in all probability the edifice would not have stood for any length of time.

Much of the thrust of a dome may be counteracted by means of an iron belt placed round it at the point where the thrust is greatest. This point I have shown to be at 70° from the crown, or 20° from the springing.

June 7, 1866.

The Annual Meeting for the election of Fellows was held this day.

Lient.-General SABINE, President, in the Chair.

The Statutes relating to the Election of Fellows having been read, Professor Brayley and Mr. Toynbee were, with the consent of the Society, nominated Scrutators to assist the Secretaries in examining the Lists.

The votes of the Fellows present having been collected, the following Candidates were declared to be duly elected into the Society :—

John Charles Bucknill, M.D.
 Rev. Frederick William Farrar.
 William Augustus Guy, M.B.
 James Hector, M.D.
 John William Kaye, Esq.
 Hugo Müller, Ph.D.
 Charles Murchison, M.D.
 William Henry Perkin, Esq.

The Ven. John Henry Pratt, M.A.
 Capt. George Henry Richards, R.N.
 Thomas Richardson, Esq., M.A.
 William Henry Leighton Russell,
 Esq.
 Rev. William Selwyn, D.D.
 Rev. Richard Townsend, M.A.
 Henry Watts, B.A.

June 14, 1866.

Lieut.-General SABINE, President, in the Chair.

The Rev. F. W. Farrar, Dr. Charles Murchison, Captain Richards, R.N., Mr. W. H. L. Russell, and Mr. Henry Watts, were admitted into the Society.

Pursuant to notice given at the last Ordinary Meeting, Franz Cornelius Donders, George Friedrich Bernhard Riemann, and Gustav Rose, were balloted for and elected Foreign Members of the Society.

The following papers were read :—

- I. "On the Anatomy of the Fovea centralis of the Human Retina." By J. W. HULKE, F.R.C.S., Assistant Surgeon to the Middlesex and Royal London Ophthalmic Hospitals. Communicated by WM. BOWMAN, Esq. Received May 26, 1866.

(Abstract).

1. The Fovea centralis is a minute circular pit in the inner surface of the retina, made by the radial divergence of the cone-fibres from a central point, by the thinning and the outward curving of the inner retinal layers towards this point, and by the peripheral location of the outer granules belonging to the central cones.

2. The inner surface of the retina declines in a rapid uniform curve from the edge to the centre of the fovea, and very gradually from the edge towards the ora retinæ; so that the edge of the fovea is the most raised part in the macula lutea, where the retina is thickest, and the centre of the fovea the most depressed part in the macula, where the retina is thinnest.

3. At the centre of the fovea, proceeding from the outer to the inner surface of the retina, we meet with the following structures in succession :—the bacillary layer and the outer limiting membrane, a small quantity of finely areolated connective tissue, the inner granule-layer and the ganglionic layer very attenuated, a thin granular band containing optic nerve-fibres, and the membrana limitans interna.